

| | Weeks | Light per day (hours) | COCO ml per 1 liters | RHIZO TONIC ml per 1 liters | CANNA ZYM ml per 1 liters | PK 13/14 ml per 1 liters | CANNA BOOST ml per 1 liters | EC+ | EC Total |
|-----------|--|-----------------------|----------------------|-----------------------------|---------------------------|--------------------------|-----------------------------|-----|----------|
| Growth | Start / rooting (3–5 days) Make substrate wet | <1 | 2.3(A) 2.3(B) | 4 | - | - | - | 1.1 | 1.3 |
| | Vegetative phase I Plant develops in volume | 0-3 ^a | 2.7(A) 2.7(B) | 2 | 2.5 | - | - | 1.3 | 1.5 |
| | Vegetative phase II Up to growth stagnation after fructification or appearance of the formation of flowers | 2-4 ^b | 3.1(A) 3.1(B) | 2 | 2.5 | - | 2 - 4 ^e | 1.5 | 1.7 |
| Flowering | Generative Period I Flowers or fruits develop in length. Growth in height achieved | 2-3 | 3.8(A) 3.8(B) | 0.5 | 2.5 | - | 2 - 4 | 1.8 | 2 |
| | Generative Period II Development of the volume (breadth) of flowers or fruit | 1 | 4.2(A) 4.2(B) | 0.5 | 2.5 | 1.5 | 2 - 4 | 2 | 2.2 |
| | Generative Period III Development of the mass (weight) of flowers or fruit | 2-3 | 2.9(A) 2.9(B) | 0.5 | 2.5 | - | 2 - 4 | 1.4 | 1.6 |
| | Generative Period IV Flowers or fruit ripening process | 1-2 | 10-12 ^c | - | - | 2.5 - 5 ^d | 2 - 4 | - | 0.2 |

Product line: COCO
 Your tanksize: 1 liters
 Scheduletype: Heavy feeding
 Water EC: 0.2
 Dosage units: ml/1 liters
 Recommended pH: 5.5 - 6.2



The guidelines in the table aren't an iron law, but can help novice growers to develop a sophisticated fertilisation strategy. The optimum fertilisation strategy is further determined by factors such as: temperature, humidity, plant species, root volume, moisture percentage in substrate, water dosage strategy, etc.

- ^a. This period varies depending on the species and number of plants per m².
Mother plants remain in this phase until the end (6 – 12 months).
 - ^b. The changeover from 18 to 12 hours varies depending on the variety. The rule of thumb is to change after 2 weeks.
 - ^c. Reduce hours of light if ripening goes too fast. Watch out for increasing Relative Humidity.
 - ^d. Double CANNAZYM dosage to 50 ml/10 liters, if substrate is reused.
 - ^e. 20 ml/ 10 liters standard. Increase to a maximum of 40 ml/10 liters for extra flowering power.
- EC:
 - EC+ value is based in mS/cm when EC water = 0.0 at 25°C, pH 6.0
 - pH:
 - Recommended pH is between 5.5 - 6.2
 - Adding pH- can increase EC
 - Use pH- grow in the vegetative as in the generative phase to lower the pH