

12 hrs 18 hrs	E	WKI	0.75ml	Imi		0.5ml	0.5ml	0.5ml	Imi	0.5ml		5.7-6.2	1.1
	VEGETATIC	WK 2	1ml	1.25ml		0.5ml	0.5ml	1ml	1ml	0.5ml		5.7-6.2	1.3
	VEC	WK 3	1ml	1.5ml		0.5ml	0.5ml	1ml	1ml	0.5ml		5.7-6.2	1.4
		WK 1	1.5ml		1.25ml	0.5ml	0.75ml	1.25ml	1ml	0.5ml	1ml	5.7-6.2	1.6
	FLOWERING	WK 2	1.5ml		1.25ml	0.5ml	0.75ml	1.5ml	1ml	0.5ml	1ml	5.7-6.2	1.6
		WK 3	1.25ml		1.7ml		1ml	1ml	1ml	0.5ml	1.5ml	5.7-6.2	1.6
		WK 4	1.5ml		2ml		1ml	1ml	1ml	0.5ml	2ml	5.7-6.2	1.8
		WK 5	1.5ml		2ml		1ml	1ml	1ml	0.5ml	2ml	5.7-6.2	1.8
		WK 6	1.5ml		2ml		1.5ml	1ml	1ml	0.5ml	2.5ml	5.7-6.2	1.9
		WK 7	1.5ml		2ml		1.5ml	1ml	1ml	0.5ml	2.5ml	5.7-6.2	1.9
		WK 8	1.2ml		1.7ml		1.5ml	1ml	1ml	0.5ml	2ml	5.7-6.2	1.7
		WK 9	Clearex	2ml									

****EC Values are guideline only as dependant an background value and size of plant.**

FEEDSHEET TIPS

- 1. When mixing Kind nutrients, add Kind Base first, then add Kind Grow or Bloom. Rinse measuring cups between doses to prevent contamination.
- 2. In warmer environments, plants will lose more water through transpiration. Applying a more dilute nutrient solution when temperatures are warmer will prevent over-fertilization.
- 3. If growing in coconut coir-based media or if using reverse osmosis water, add 0.75ml-1.25ml of Botanicare Cal-Mag Plus to the nutrient solution recipe.
- 4. Maintain a nutrient solution pH range between 5.7 6.2.
- 5. The optimal temperature range of the nutrient solution is... 18 21°C.
- 6. Use nutrient solution immediately after mixing or keep solution circulating to prevent settling.
- 7. If using a recirculating system, maintain water level in reservoir by adding fresh water and nutrients as needed. Change solution every week.
- 8. Additional Botanicare supplements have been scientifically formulated to meet plant needs during important phases of growth and development. Use Hydroguard for maximum root protection and Clearex to break ionic bonds in the grow media while flushing or in the case of over fertilization.